

On some definite integrals connecting with certain infinite series

Ramesh Kumar Muthumalai *

May 6, 2010

Abstract. We show some definite integrals connecting to infinite series, studied in Ramanujan's paper, titled "On question 330 of Professor Sanjana". We present few recursive methods to evaluate these definite integrals in various cases and we generalize this, to evaluate simliar kind of integrals through infinite series.

AMS subject classification: 33E20; 33E50; 40A05.

Keywords: Definite integrals; Question 330 of Professor Sanjana; Ramanujan papers; Infinite series.

1. Introduction The classical table of integrals by I.S. Gradshteyn and I.M. Ryzhik [2] contains many entries related to definite integrals in the combinations of powers and algebraic functions of exponentials. In particular, the following integral is given for some special cases of a and α

$$\int_0^\infty x^\alpha e^{-bx} (1 - e^{-x})^a dx \quad (1.1)$$

In this paper, we connect this definite integral to infinite series given in [3]. We evaluate them for various cases a and α by using recursive methods. Further, we have shown evaluation of combination of trigonometric functions and powers. Finally, we present summation of the series of this type

$$\frac{1}{b^n} - \frac{a}{1!(b+1)^n} \frac{1}{a} + \frac{a(a-1)}{2!(b+2)^n} \left(\frac{1}{a} + \frac{1}{a-1} \right) - \dots \quad (1.2)$$

*Department of Mathematics, Sindhi college of arts and science, chennai-77, Tamil Nadu, India. Email id: ramjan_80@yahoo.com. Home page URL: http://ramjan_07.page.tl/.

2. Definite integrals connecting with infinite series. Consider the following definite integrals for $b > 0$ and $\alpha \geq 0$

$$\begin{aligned}
& \int_0^\infty x^\alpha e^{-bx} (1 - e^{-x})^a dx \\
&= \int_0^\infty x^\alpha e^{-bx} \left(1 - \frac{a}{1!} e^{-x} + \frac{a(a-1)}{2!} e^{-2x} - \dots \right) dx \\
&= \Gamma(\alpha + 1) \left(\frac{1}{b^{\alpha+1}} - \frac{a}{1!(b+1)^{\alpha+1}} + \frac{a(a-1)}{2!(b+2)^{\alpha+1}} - \dots \right) \\
&= \Gamma(\alpha + 1) \phi(a, b, \alpha)
\end{aligned}$$

Where

$$\phi(a, b, \alpha) = \frac{1}{b^{\alpha+1}} - \frac{a}{1!(b+1)^{\alpha+1}} + \frac{a(a-1)}{2!(b+2)^{\alpha+1}} - \dots \quad (2.1)$$

Thus, we obtain

$$\int_0^\infty x^\alpha e^{-bx} (1 - e^{-x})^a dx = \Gamma(\alpha + 1) \phi(a, b, \alpha) \quad (2.2)$$

If $\alpha = n, n \in \mathbb{N}$ then

$$\int_0^\infty x^n e^{-bx} (1 - e^{-x})^a dx = n! \phi(a, b, n) \quad (2.3)$$

Also, (2.3) can be written as

$$\int_0^1 \log^n(1-t) (1-t)^{b-1} t^a dt = (-1)^n n! \phi(a, b, n) \quad (2.4)$$

Differentiating (2.3) m times with respect to a , we find that

$$\int_0^\infty x^n e^{-bx} (1 - e^{-x})^a \log^m(1 - e^{-x}) dx = n! \phi_a^{(m)}(a, b, n) \quad (2.5)$$

The above integral can be written as

$$\int_0^1 \log^n(1-t) \log^m t (1-t)^{b-1} t^a dt = (-1)^n n! \phi_a^{(m)}(a, b, n) \quad (2.6)$$

Interchanging m and n in (2.6), we find

$$(-1)^m m! \phi_b^{(n)}(b-1, a+1, m) = (-1)^n n! \phi_a^{(m)}(a, b, n) \quad (2.7)$$

If $n = 0$

$$(-1)^m m! \phi(b-1, a+1, m) = \phi_a^{(m)}(a, b, 0) \quad (2.8)$$

Thus, through the infinite series $\phi(a, b, 0)$, we can evaluate four different form of integrals. Similiarly, if we start from the infinite series for $b > 0$ and $\alpha \geq 0$

$$\frac{1}{b^{\alpha+1}} + \frac{a}{1!(b+1)^{\alpha+1}} + \frac{a(a-1)}{2!(b+2)^{\alpha+1}} - \dots = \tilde{\phi}(a, b, \alpha) \quad (2.9)$$

then, we obtain

$$\int_0^\infty x^\alpha e^{-bx} (1 + e^{-x})^a dx = \Gamma(\alpha + 1) \tilde{\phi}(a, b, \alpha) \quad (2.10)$$

Generalizing equations (2.3) and (2.10), for some $|\beta| < 1$

$$\int_0^\infty x^\alpha e^{-bx} (1 + \beta e^{-x})^a dx = \Gamma(\alpha + 1) \Psi(a, b, \beta, \alpha) \quad (2.11)$$

Where

$$\frac{1}{b^{\alpha+1}} + \frac{a}{1!(b+1)^{\alpha+1}} \beta + \frac{a(a-1)}{2!(b+2)^{\alpha+1}} \beta^2 + \dots = \Psi(a, b, \beta, \alpha) \quad (2.12)$$

Again, consider the following definite integral for $b > 0$ and $\alpha \geq 0$

$$\begin{aligned} \int_0^\infty x^\alpha e^{-ibx} (1 - e^{-ix})^a dx \\ &= \int_0^\infty x^\alpha e^{-ibx} \left(1 - \frac{a}{1!} e^{-ix} + \frac{a(a-1)}{2!} e^{-2ix} - \dots \right) dx \\ &= \frac{\Gamma(\alpha + 1)}{i^{\alpha+1}} \left(\frac{1}{b^{\alpha+1}} - \frac{a}{1!(b+1)^{\alpha+1}} + \frac{a(a-1)}{2!(b+2)^{\alpha+1}} - \dots \right) \\ &= -\Gamma(\alpha + 1) \phi(a, b, \alpha) \left(\sin \frac{\alpha\pi}{2} + i \cos \frac{\alpha\pi}{2} \right) \end{aligned} \quad (2.13a)$$

On the other hand, we have

$$\begin{aligned} \int_0^\infty x^\alpha e^{-ibx} (1 - e^{-ix})^a dx = \\ 2^a \int_0^\infty x^\alpha \sin^\alpha \frac{x}{2} \left(\cos \left[\frac{\pi-x}{2} a + bx \right] - i \sin \left[\frac{\pi-x}{2} a + bx \right] \right) dx \end{aligned} \quad (2.13b)$$

Comparing (2.13a) and (2.13b), we obtain

$$\int_0^\infty x^\alpha \sin^a \frac{x}{2} \cos \left(\frac{\pi-x}{2} a + bx \right) dx = -2^{-a} \Gamma(\alpha + 1) \phi(a, b, \alpha) \sin \frac{\alpha\pi}{2} \quad (2.14)$$

$$\int_0^\infty x^\alpha \sin^a \frac{x}{2} \sin \left(\frac{\pi-x}{2} a + bx \right) dx = 2^{-a} \Gamma(\alpha + 1) \phi(a, b, \alpha) \cos \frac{\alpha\pi}{2} \quad (2.15)$$

Expanding the cosine function in the integral (2.14),

$$\begin{aligned} \int_0^\infty x^\alpha \sin^a \frac{x}{2} \left(\cos \frac{\pi a}{2} \cos \left(b - \frac{a}{2} \right) x - \sin \frac{\pi a}{2} \sin \left(b - \frac{a}{2} \right) x \right) \\ = -2^{-a-\alpha-1} \Gamma(\alpha+1) \phi(a, b, \alpha) \sin \frac{\alpha\pi}{2} \end{aligned}$$

Setting $u = 2b - a$, then (2.14) gives

$$\begin{aligned} \int_0^\infty x^\alpha \sin^a x \left(\cos \frac{\pi a}{2} \cos ux - \sin \frac{\pi a}{2} \sin ux \right) \\ = -2^{-a-\alpha-1} \Gamma(\alpha+1) \phi \left(a, \frac{u+a}{2}, \alpha \right) \sin \frac{\alpha\pi}{2} \end{aligned}$$

Setting $\lambda_c = \int_0^\infty x^\alpha \sin^a x \cos uxdx$ and $\lambda_s = \int_0^\infty x^\alpha \sin^a x \sin uxdx$

$$\cos \frac{\pi a}{2} \lambda_c - \sin \frac{\pi a}{2} \lambda_s = -2^{-a-\alpha-1} \Gamma(\alpha+1) \phi \left(a, \frac{u+a}{2}, \alpha \right) \sin \frac{\alpha\pi}{2} \quad (2.16a)$$

Similiarly from (2.15), we have

$$\sin \frac{\pi a}{2} \lambda_c + \cos \frac{\pi a}{2} \lambda_s = 2^{-a-\alpha-1} \Gamma(\alpha+1) \phi \left(a, \frac{u+a}{2}, \alpha \right) \cos \frac{\alpha\pi}{2} \quad (2.16b)$$

Solving (2.16a) and (2.16b), we get

$$\lambda_c = 2^{-a-\alpha-1} \Gamma(\alpha+1) \phi \left(a, \frac{u+a}{2}, \alpha \right) \sin(a-\alpha) \frac{\pi}{2} \quad (2.17)$$

$$\lambda_s = 2^{-a-\alpha-1} \Gamma(\alpha+1) \phi \left(a, \frac{u+a}{2}, \alpha \right) \cos(a-\alpha) \frac{\pi}{2} \quad (2.18)$$

Differentiating (2.17) and (2.18) m times, with respect to a , we can evaluate the following definite integrals respectively

$$\int_0^\infty x^\alpha \log^m \sin x \sin^a x \cos uxdx \quad \text{and} \quad \int_0^\infty x^\alpha \log^m \sin x \sin^a x \sin uxdx$$

Similiarly, if we start from the integral $\int_0^\infty x^\alpha e^{-ibx} (1 + e^{-ix})^a dx$ we find

$$\int_0^\infty x^\alpha \cos^a \frac{x}{2} \cos \left(\frac{a}{2} + b \right) x dx = -2^{-a} \Gamma(\alpha+1) \tilde{\phi}(a, b, \alpha) \sin \frac{\alpha\pi}{2}$$

$$\int_0^\infty x^\alpha \cos^a \frac{x}{2} \sin \left(\frac{a}{2} + b \right) x dx = 2^{-a} \Gamma(\alpha+1) \tilde{\phi}(a, b, \alpha) \cos \frac{\alpha\pi}{2}$$

Setting $v = a + 2b$ and after simplification, we find

$$\int_0^\infty x^\alpha \cos^a x \cos vxdx = -2^{-a-\alpha-1} \Gamma(\alpha+1) \tilde{\phi} \left(a, \frac{v-a}{2}, \alpha \right) \sin \frac{\alpha\pi}{2} \quad (2.19)$$

$$\int_0^\infty x^\alpha \cos^a x \sin vxdx = 2^{-a-\alpha-1} \Gamma(\alpha+1) \tilde{\phi} \left(a, \frac{v-a}{2}, \alpha \right) \cos \frac{\alpha\pi}{2} \quad (2.20)$$

3. Algebraic recursive method Let we start by deriving a recursive method connecting algebraic functions to infinite series, for some $t < 1$, we know that from binomial theorem [1]

$$b - \frac{p}{1!}(b+1)t + \frac{p(p-1)}{2!}(b+2)t^2 - \dots = b(1-t)^p - pt(1-t)^{p-1}$$

Similiarly, we can write

$$b^2 - \frac{p}{1!}(b+1)^2t + \frac{p(p-1)}{2!}(b+2)^2t^2 - \dots = b^2(1-t)^p - (2b+1)pt(1-t)^{p-1} + p(p-1)t^2(1-t)^{p-2}$$

Generalize this to positive integer m , we have

$$\sum_{i=0}^{\infty} (-1)^i \binom{p}{i} (b+i)^{m-1} t^i = \sum_{k=1}^m A_k^{(m)} t^{k-1} (1-t)^{p-k+1} \quad (3.1)$$

Multiplying by t^b on (3.1)

$$\sum_{i=0}^{\infty} (-1)^i \binom{p}{i} (b+i)^{m-1} t^{i+b} = \sum_{k=1}^m A_k^{(m)} t^{k+b-1} (1-t)^{p-k+1}$$

Differentiating with respect to t ,

$$\begin{aligned} & \sum_{i=0}^{\infty} (-1)^i \binom{p}{i} (b+i)^m t^{i+b-1} \\ &= \sum_{k=1}^m A_k^{(m)} ((k+b-1)t^{k+b-2}(1-t)^{p-k+1} - (p-k+1)t^{k+b-1}(1-t)^{p-k}) \end{aligned}$$

Now rearranging

$$\begin{aligned} &= A_1^{(m)} b t^{b-1} (1-t)^p \\ &+ \sum_{k=2}^m \left(-(p-k+2)A_{k-1}^{(m)} + (b+k-1)A_k^{(m)} t^{b+k-2}(1-t)^{p-k+1} \right) \\ &- (p-m+1)A_m^{(m)} t^{m+b-1}(1-t)^{p-m} \end{aligned}$$

Cancelling t^{b-1} on both sides

$$\begin{aligned} & A_1^{(m)} b (1-t)^p \\ &+ \sum_{k=2}^m \left(-(p-k+2)A_{k-1}^{(m)} + (b+k-1)A_k^{(m)} t^{k-1}(1-t)^{p-k+1} \right) \\ &- (p-m+1)A_m^{(m)} t^m (1-t)^{p-m} = \sum_{i=0}^{\infty} (-1)^i \binom{p}{i} (b+i)^m t^i \quad (3.2) \end{aligned}$$

Now (3.2) can be written in the form of equation (3.1). If we replace m by $m + 1$

$$\sum_{k=1}^{m+1} A_k^{(m+1)} t^{k-1} (1-t)^{p-k+1} = \sum_{i=0}^{\infty} (-1)^i \binom{p}{i} (b+i)^m t^i \quad (3.3)$$

Comparing (3.2) and (3.3)

$$\begin{aligned} A_1^{(m+1)} &= b A_1^{(m)} \\ A_k^{(m+1)} &= -(p-k+2) A_{k-1}^{m+1} + (b+k-1) A_k^m, \quad k = 2, 3, \dots, m \\ A_{m+1}^{(m+1)} &= -(p-m+1) A_m^{(m)} \end{aligned} \quad (3.4)$$

If we put $m = 0$ in (3.1) then we find that $A_1^{(1)} = 1$. So, we can evaluate all A 's recursively through equation (3.4).

4. Evaluation of definite integrals through infinite series. In section 2, we show some definite integrals through infinite series. Here, we evaluate them for various cases.

Evaluation of $\phi(a, b, \alpha)$

This integral exists, if $\alpha \geq 0$ and $b > 0$. It is clear that, from equation (2.2)

$$\int_0^{\infty} x^{\alpha} e^{-bx} (1 - e^{-x})^a dx = \Gamma(\alpha + 1) \phi(a, b, \alpha)$$

Now, we evaluate them for various cases of α and a through infinite series $\phi(a, b, \alpha)$.

Case 1. If $\alpha = n$ is an integer, then we can use Ramanujan formula, to evaluate infinite series $\phi(a, b, \alpha)$ as follow as

$$n \phi(a, b, n) = \sigma_1 \phi(a, b, n-1) + \sigma_2 \phi(a, b, n-2) + \dots + \sigma_n \phi(a, b, 0) \quad (4.1)$$

Where

$$\sigma_k = \frac{1}{b^k} - \frac{1}{(a+b+1)^k} + \frac{1}{(b+1)^k} - \frac{1}{(a+b+2)^k} + \dots \quad (4.2)$$

and $k = 1, 2, 3, \dots$

Example. If we take $a = -\frac{1}{2}$, $b = \frac{1}{4}$ and $n = 0, 1, 2$, then $\phi(-\frac{1}{2}, \frac{1}{4}, n)$ can be calculated from (4.1). Finally, using (2.2) we find

$$\int_0^{\infty} x e^{-\frac{x}{4}} (1 - e^{-x})^{-\frac{1}{2}} dx = \frac{\Gamma(\frac{1}{4})^2 \pi}{\sqrt{2\pi}}$$

$$\int_0^\infty x^2 e^{-\frac{x}{4}} (1 - e^{-x})^{-\frac{1}{2}} dx = \frac{\Gamma(\frac{1}{4})^2}{\sqrt{2\pi}} (\pi^2 + 16S'_2)$$

$$\int_0^\infty x^3 e^{-\frac{x}{4}} (1 - e^{-x})^{-\frac{1}{2}} dx = \frac{\Gamma(\frac{1}{4})^2}{\sqrt{2\pi}} (5\pi^3 + 48S'_2 + 128S'_3)$$

Where

$$S'_r = \frac{1}{1^r} - \frac{1}{3^r} + \frac{1}{5^r} - \frac{1}{7^r} + \dots + \infty$$

Case 2. If $\alpha = \mu + m, \mu > 0, m \in N$ and $a = -1, -2, -3, \dots$ then putting $p = -1, t = e^{-x}$ in (3.3) and multiplying by $x^{\mu+m} e^{-bx}$ on both sides, we find

$$\sum_{k=1}^{m+1} A_k^{(m+1)} \frac{x^{\mu+m} e^{-x(b+k-1)}}{(1 - e^{-x})^k} = \sum_{i=0}^{\infty} (b+i)^m x^{\mu+m} e^{-(b+i)x}$$

Integrating on $(0, \infty)$,

$$\sum_{k=1}^{m+1} A_k^{(m+1)} \int_0^\infty \frac{x^{\mu+m} e^{-x(b+k-1)}}{(1 - e^{-x})^k} dx = \sum_{i=0}^{\infty} (b+i)^m \int_0^\infty x^{\mu+m} e^{-(b+i)x} dx \quad (4.3)$$

Using (2.2) and after simplification, we get

$$\sum_{k=1}^{m+1} A_k^{(m+1)} \phi(-k, b+k-1, \mu+m) = \zeta(\mu+1, b) \quad (4.4)$$

Example. If we put $m = 0$ and $\mu > 0$ in (4.4)

$$\phi(-1, b, \mu) = \zeta(\mu+1, b)$$

Putting $m = 1$ in (4.4)

$$\phi(-2, b+1, \mu+1) = \zeta(\mu+1, b) - b\zeta(\mu+2, b)$$

Putting $m = 2$ in (4.4)

$$\phi(-3, b+2, \mu+2) = \zeta(\mu+1, b) - (2b+1)\zeta(\mu+2, b) + b(b+1)\zeta(\mu+3, b)$$

Case 3. If $\alpha = \mu + m, \mu > 0, m \in N$ and $a = \gamma + m \neq -1, -2, -3, \dots$ then putting $\gamma + m = p, t = e^{-x}$ in (3.3) and multiplying by $x^{\mu+m} e^{-bx}$ on both sides and integrating on $(0, \infty)$

$$\sum_{k=1}^{m+1} A_k^{(m+1)} \int_0^\infty x^{\mu+m} e^{x(-b-k+1)} (1 - e^{-x})^{\gamma+m-k+1} dx$$

$$= \sum_{i=0}^{\infty} (-1)^i \binom{\gamma+m}{i} (b+i)^m \int_0^\infty x^{\mu+m} e^{-(b+i)x} dx$$

Using (2.2) and after simplification, we get

$$\sum_{k=1}^{m+1} A_k^{(m+1)} \phi(\gamma + m - k + 1, b + k - 1, \mu + m) = \phi(\gamma + m, b, \mu) \quad (4.5)$$

Example. If we take $\gamma = -\frac{1}{2}$, $b = \frac{1}{4}$, $\mu = 0$ and $m = 1$ in (3.3), then

$$\frac{1}{4} \phi\left(-\frac{1}{2}, \frac{1}{4}, 1\right) + \frac{1}{2} \phi\left(-\frac{3}{2}, \frac{5}{4}, 1\right) = \phi\left(-\frac{1}{2}, \frac{1}{4}, 0\right)$$

So that

$$\int_0^\infty x e^{-\frac{5x}{4}} (1 - e^{-x})^{-\frac{3}{2}} dx = 2 \frac{\Gamma(\frac{1}{4})^2}{\sqrt{2\pi}} \left(1 - \frac{\pi}{4}\right)$$

Evaluation of $\tilde{\phi}(a, b, \alpha)$

This integral exists, if $\alpha \geq 0$ and $b > 0$. From the equation (2.10). It is clear that

$$\int_0^\infty x^\alpha e^{-bx} (1 + e^{-x})^a dx = \Gamma(\alpha + 1) \tilde{\phi}(a, b, \alpha)$$

Now, we evaluate them for various cases of $\phi(a, b, \alpha)$.

Case 1. If $\alpha = \mu + m$, $\mu \geq 0$, $m \in N$ and $a = -1, -2, -3, \dots$

If $a = 1$, then

$$\int_0^\infty \frac{x^\alpha e^{-bx}}{1 - e^{-2x}} dx = \frac{\Gamma(\alpha + 1)}{2^\alpha} \phi\left(-1, \frac{b}{2}, \alpha\right)$$

Also,

$$\begin{aligned} \int_0^\infty \frac{x^\alpha e^{-bx}}{1 + e^{-x}} dx &= \frac{\Gamma(\alpha + 1)}{2^\alpha} \phi\left(-1, \frac{b}{2}, \alpha\right) - \Gamma(\alpha + 1) \phi(-1, b, \alpha) \\ \tilde{\phi}(-1, b, \alpha) &= \frac{1}{2^\alpha} \phi\left(-1, \frac{b}{2}, \alpha\right) - \phi(-1, b, \alpha) \end{aligned} \quad (4.6)$$

Putting $p = -1$, $t = e^{-x}$ in (3.3), multiplying by $x^{\mu+m} e^{-bx}$ on both sides and integrating $(0, \infty)$, we get

$$\sum_{k=1}^{m+1} A_k^{(m+1)} (-1)^{k-1} \tilde{\phi}(-k, b + k - 1, \mu + m) = \sum_{i=0}^\infty \frac{(-1)^i}{(b + i)^{\mu+1}} \quad (4.7)$$

Case 2. If $\alpha = \mu + m$, $\mu > 0$ and $a = \gamma + m \neq -1, -2, -3, \dots$ Similiarly, putting $p = \gamma + m$, $t = -e^{-x}$ in (3.5), multiplying by $x^{\mu+m} e^{-bx}$ on both sides

and integrating on $(0, \infty)$

$$\sum_{k=1}^{m+1} (-1)^{k-1} A_k^{(m+1)} \tilde{\phi}(\gamma + m - k + 1, b + k - 1, \mu + m) = \tilde{\phi}(\gamma + m, b, \mu) \quad (4.8)$$

Evaluation of $\Psi(a, b, \beta, \alpha)$

Putting $p = \gamma + m, t = -\beta e^{-x}, |\beta| < 1$ in (3.3), multiplying by $x^{\mu+m} e^{-bx}$ and integrating on $(0, \infty)$, then

$$\sum_{k=1}^{m+1} A_k^{(m+1)} (-\beta)^{k-1} \Psi(\gamma + m - k + 1, b + k - 1, \beta, \mu + m) = \Psi(\gamma + m, b, \beta, \mu) \quad (4.9)$$

Example. If we put $m = 0$ and $\mu \geq 0$ in (4.9)

$$\Psi(-1, b, \beta, \mu) = \Phi(\beta, \mu + 1, b)$$

Putting $m = 1$ in (4.9)

$$\Psi(-2, b + 1, \beta, \mu + 1) = \frac{1}{\beta} (\Phi(\beta, \mu + 1, b) - b\Phi(\beta, \mu + 2, b))$$

Where $\Phi(\beta, \mu + 1, b)$ is the Lerch function. To show an another interesting example, consider the following definite integral from [2], pp 350.

$$\int_{-\infty}^{\infty} \frac{x^2 e^{-bx}}{1 + e^{-x}} dx = \pi^3 \csc b\pi (2 - \sin^2 b\pi)$$

Now, for $p = -1, -2, -3, \dots, t = -\beta e^{-x}$ in (3.5), multiply and divide by $x^2 e^{-bx}$ and $1 + \beta e^{-x}$ respectively, then, after integrating on $(-\infty, \infty)$

$$\begin{aligned} \sum_{k=1}^{m+1} A_k^{(m+1)} \int_{-\infty}^{\infty} \frac{(-\beta)^{k-1} x^2 e^{-x(b+k-1)}}{(1 + e^{-x})(1 + \beta e^{-x})^k} dx &= \sum_{i=0}^{\infty} \beta^i (b + i)^m \int_{-\infty}^{\infty} \frac{x^2 e^{-(b+i)x}}{1 + e^{-x}} dx \\ &= \pi^3 \csc b\pi (2 - \sin^2 b\pi) \sum_{i=0}^{\infty} \beta^i (b + i)^m \end{aligned}$$

If we put $m = 0$, we obtain

$$\int_{-\infty}^{\infty} \frac{x^2 e^{-bx}}{(1 + e^{-x})(1 + \beta e^{-x})} dx = \frac{\pi^3}{1 - \beta} \csc b\pi (2 - \sin^2 b\pi)$$

By putting $m = 1, 2, \dots$, we can easily evaluate definite integrals of the type $\int_{-\infty}^{\infty} \frac{x^2 e^{-bx}}{(1 + e^{-x})(1 + \beta e^{-x})^{m+1}} dx$. Thus, using new algebraic recursive method, we can evaluate similar type of integrals.

5. Summation of Infinite series. To evaluate integrals that given in equation (2.6) and (2.10a). We need to know summation of the series of the type (1.2). It can be found by differentiating Ramanujan recursive formula (2.4) successively with respect to a . but, it may requires some additional calculation and manipulation time. Here, we present alternative formula that require slight lesser computation then the differentiation of (2.4). We derive this formula almost similar way of Ramanujan formula. For $a \neq -1, -2, -3, \dots$, using Taylor series [1] near $p = 0$

$$\log \Gamma(p+a+1) = \log \Gamma(a+1) + \psi(a+1)\frac{p}{1} + \zeta(2, a+1)\frac{p^2}{2} - \zeta(3, a+1)\frac{p^3}{3} + \dots \quad (5.1)$$

Where

$$\psi(a+1) = \lim_{n \rightarrow \infty} \log n - \left(\frac{1}{a+1} + \frac{1}{a+2} + \frac{1}{a+3} + \dots + \frac{1}{a+n} \right) \quad (5.2)$$

and

$$\zeta(r, a+1) = \frac{1}{(a+1)^r} + \frac{1}{(a+2)^r} + \frac{1}{(a+3)^r} + \dots \quad (5.3)$$

Let us take

$$f(p, a, b) = \int_0^1 x^{p+b-1} (1-x)^a dx$$

Expanding in ascending powers of p and integrating by x ,

$$= \frac{1}{b+p} - \frac{a}{1!(b+p+1)} + \frac{a(a-1)}{2!(b+p+2)} - \frac{a(a-1)(a-2)}{3!(b+p+3)} + \dots$$

$$f(p, a, b) = \phi(a, b, 0) - p\phi(a, b, 1) + p^2\phi(a, b, 2) - p^3\phi(a, b, 3) + \dots \quad (5.4)$$

Now, differentiating with respect to a ,

$$f'(p, a, b) = \phi'_a(a, b, 0) - p\phi'_a(a, b, 1) + p^2\phi'_a(a, b, 2) + \dots \quad (5.4a)$$

$$f(p, a, b) = \frac{\Gamma(p+b)\Gamma(a+1)}{\Gamma(p+a+b+1)}$$

Therefore,

$$\log f(p, a, b) = \log \Gamma(p+b) + \log \Gamma(a+1) - \log \Gamma(p+a+b+1)$$

Again, differentiating with respect to a

$$\frac{f'(p, a, b)}{f(p, a, b)} = \psi(a+1) - \psi(p+a+b+1)$$

$$\frac{f'(p, a, b)}{f(p, a, b)} = \psi(a+1) - (\psi(a+b+1) + p\zeta(2, a+b+1) - p^2\zeta(3, a+b+1) + \dots)$$

Using (5.4) and (5.4a)

$$\begin{aligned} & \phi'_a(a, b, 0) - p\phi'_a(a, b, 1) + p^2\phi'_a(a, b, 2) - \dots \\ &= (\psi(a+1) - \psi(a+b+1) - p\zeta(2, a+b+1) + p^2\zeta(3, a+b+1) - \dots) \times \\ & \quad (\phi(a, b, 0) - p\phi(a, b, 1) + p^2\phi(a, b, 2) - p^3\phi(a, b, 3) + \dots) \end{aligned}$$

Equating coefficients of p

$$\begin{aligned} \phi'_a(a, b, 0) &= (\psi(a+1) - \psi(a+b+1))\phi(a, b, 0) \\ \phi'_a(a, b, 1) &= (\psi(a+1) - \psi(a+b+1))\phi(a, b, 1) + \zeta(2, a+b+1)\phi(a, b, 0) \\ \phi'_a(a, b, 2) &= (\psi(a+1) - \psi(a+b+1))\phi(a, b, 2) \\ & \quad + \zeta(2, a+b+1)\phi(a, b, 1) + \zeta(3, a+b+1)\phi(a, b, 0) \end{aligned}$$

Generalize this, we obtain

$$\begin{aligned} \phi'_a(a, b, n) &= (\psi(a+1) - \psi(a+b+1))\phi(a, b, n) \\ & \quad + \zeta(2, a+b+1)\phi(a, b, n-1) + \zeta(3, a+b+1)\phi(a, b, n-2) \\ & \quad + \dots + \zeta(n+1, a+b+1)\phi(a, b, 0) \end{aligned} \quad (5.5)$$

Further $\phi''(a, b, n), \phi'''(a, b, n), \dots$ can be found by successive differentiation of (5.5). Also, summation of the series (1.2) is given by

$$\begin{aligned} & \frac{1}{b^n} - \frac{a}{1!(b+1)^n} \frac{1}{a} + \frac{a(a-1)}{2!(b+2)^n} \left(\frac{1}{a} + \frac{1}{a-1} \right) - \dots \\ & \quad = \frac{1}{b^n} + \phi'_a(a, b, n-1) \end{aligned} \quad (5.6)$$

Simplifying (5.6) and letting a tends to 0, we obtain

$$\frac{1}{1(b+1)^n} + \frac{1}{2(b+2)^n} + \frac{1}{3(b+3)^n} - \dots = \phi'_a(0, b, n-1) \quad (5.7)$$

Thus,

$$\phi'_a(0, b, n) = -\frac{C + \psi(b+1)}{b^{n+1}} + \frac{\zeta(2, b+1)}{b^n} + \dots + \frac{\zeta(n+1, b+1)}{b} \quad (5.8)$$

Where C is Euler constant.

Examples. If we take $a = -\frac{1}{2}$ and $b = \frac{1}{4}$ then

$$\begin{aligned} & \frac{1}{1^n} + \frac{1}{2.5^n} \frac{1}{1} + \frac{1.3}{2.4.9^n} \left(\frac{1}{1} + \frac{1}{3} \right) + \frac{1.3.5}{2.4.6.13^n} \left(\frac{1}{1} + \frac{1}{3} + \frac{1}{5} \right) \dots \\ & \quad = 4^n - \phi'_a\left(-\frac{1}{2}, \frac{1}{4}, n-1\right) \end{aligned}$$

If $n = 0, 1$, then $\phi'(-\frac{1}{2}, \frac{1}{4}, n)$ can be calculated from (5.5). Finally, using (2.6) we find

$$\int_0^1 \log t (1-t)^{-\frac{3}{4}} t^{-\frac{1}{2}} dt = \frac{\Gamma(\frac{1}{4})^2}{\sqrt{2\pi}} \left(-\frac{\pi}{2} + \ln 2 \right)$$

$$\int_0^1 \log(1-t) \log t (1-t)^{-\frac{3}{4}} t^{-\frac{1}{2}} dt = \frac{\Gamma(\frac{1}{4})^2}{\sqrt{2\pi}} \left(\pi \left(-\frac{\pi}{2} + \ln 2 \right) + \zeta \left(2, \frac{3}{4} \right) \right)$$

6. Conclusion. We conclude this paper, by connecting some definite integrals through infinite series given by Ramanujan [3]. Also, we studied various recursive methods to evaluate these definite integrals through Ramanujan [3] infinite series, Riemann zeta function, etc. Numerical examples are also shown to explain the procedure of these recursive methods.

References

- [1] Apostol, *Mathematical Analysis, 2 Ed*, Addison-Wesley Publishing Company(1974), USA
- [2] Gradshteyn I.S & Ryzhik I.M, *Tables of Integrals, Series and Products, 6 Ed*, Academic Press (2000), USA
- [3] Ramanujan S, *On question 330 of Professor Sanjana*, JIMS, IV (1912) India.